

Colonoscopy with CLENPIQ Prep

Colonoscopy Preparation Timeline

Check boxes as you progress!



21 Days Before or ASAP: *Purchase / Pick up items for CLENPIQ Prep*

Pick up prescription at your local pharmacy:

- Fill your prescription for CLENPIQ prep at your local pharmacy
- Please call our office at (612) 871-1145 if you did not receive a prescription.



7 Days Before:

- STOP** taking fiber and iron supplements.
- If you have diabetes, contact your monitoring provider for further direction on insulin and/or blood sugar management.



3 Days Before:

- Begin low-fiber diet (10-15g/day.) **AVOID** seeds, nuts, popcorn, corn, whole grains, quinoa, fruit, etc.

For additional details on following a low fiber diet, please visit www.mngi.com/conditions/low-fiber-diet



NEXT PAGE >



2 Days Before:

- Stay well-hydrated.
- **STOP** eating solid foods at **11:45PM** and begin clear liquid diet (only liquids you can see through).

For additional details on following a clear liquid diet, please visit <https://www.mngi.com/conditions/clear-liquid-diet>



1 Day Before:

- No solid food. Continue clear liquid diet. **NO** alcohol, red liquids, dairy, oils, pulp, etc.
- At **5PM**, drink first dose (1 bottle) of CLENPIQ.
- At your own pace **over the next 5 hours**, drink five 8 oz glasses of clear liquid.


At 5PM



Drink 1 Bottle
CLENPIQ



During next 5 hours



Drink five (8 oz)
glasses of clear liquids

Continue liquid diet ONLY



Day of Procedure:

- Continue clear liquid diet **ONLY**.
- Take your morning medications with sips of water **at least 3 hours prior** to your procedure.
 - This includes heart and blood pressure medications, blood thinners (if you have not been told to stop these by our office), methadone, and anti-seizure medications.
 - DO NOT take any chewable vitamins or supplements.
- **DO NOT** use chewing tobacco, THC, Marijuana, or illicit drugs.

NEXT PAGE





5 Hours Before:

- ❑ Drink second dose (1 bottle) of CLENPIQ.
- ❑ At your pace **over the next 2 hours**, drink three 8oz glasses of clear liquid.

5 hours before



Drink other (1) bottle
of CLENPIQ



During next 2 hours



Drink three (8 oz) glasses
of clear liquids



3 Hours Before:

- ❑ **STOP** consuming all liquids. **NOTHING BY MOUTH** until after the colonoscopy.

Ensure stool is clear liquid **ONLY** before your appointment.

If uncertain, contact our office at 612-871-1145.



Additional Information:

- **Medication Adjustments:** If taking GLP-1 agonists (Ozempic, Weygovy, Trulicity, Rybelsus, Saxenda, Byetta, or others) or SGLT-2 inhibitors (Invokana, Farxiga, Jardiance, Steglatro) follow specific holding guidelines.
<https://www.mngi.com/important-announcement-patients-having-procedure>
- **Colon Cleansing Tips:** Stay well hydrated, remain near a toilet, and continue drinking the prep solution every 15 minutes. You may want to keep the solution chilled but do not add ice. You may also consider using a straw.
- **Appointment Changes:** Reschedule if experiencing flu-like symptoms or you have newly diagnosed medical conditions.
- **Responsible Person Requirement:** Ensure you have a companion with you for check-in, discharge, and transportation.
- **How will I know I'm ready?** When you are finished with your prep you should not have any solid stool, only clear liquid stool. You should be able to see to the bottom of the toilet bowl. If you are unsure whether your colon is cleaned out, please call our office at 612-871-1145 before you leave for your appointment.
- **Questions?** For answers to frequently asked questions about your colonoscopy prep and visit. <https://www.mngi.com/colonoscopy-faqs>